



Mezgebo, G. B., Monahan, F. J., McGee, M., O'Riordan, E. G., Picard, B., Richardson, I., & Moloney, AP. (2016). Biochemical and organoleptic characteristics of muscle from early and late maturing bulls in different production systems. *animal*.  
<https://doi.org/10.1017/S175173111600272X>

Peer reviewed version

Link to published version (if available):  
[10.1017/S175173111600272X](https://doi.org/10.1017/S175173111600272X)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via CUP at <https://www.cambridge.org/core/journals/animal/article/biochemical-and-organoleptic-characteristics-of-muscle-from-early-and-late-maturing-bulls-in-different-production-systems/56D7850DC7294699228F2A35D836DB83>. Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

**Biochemical and organoleptic characteristics of muscle from early and late maturing bulls in different production systems**

G.B. Mezgebo<sup>1,2</sup>, F.J. Monahan<sup>1\*</sup>, M. McGee<sup>2</sup>, E.G. O’Riordan<sup>2</sup>, B. Picard<sup>3</sup>, R.I.

Richardson<sup>4</sup> and A.P. Moloney<sup>2</sup>

<sup>1</sup>*University College Dublin, School of Agriculture and Food Science, Dublin 4, Ireland*

<sup>2</sup>*Teagasc, Animal and Grassland Research and Innovation Centre, Grange, Dunsany, Co. Meath, Ireland*

<sup>3</sup>*Unité de Recherche sur les Herbivores, INRA Clermont-Ferrand, Theix, F-63122 Saint-Genès-Champanelle, France*

<sup>4</sup>*University of Bristol, School of Clinical Veterinary Science, Langford, Bristol BS40 5DU, England, United Kingdom*

\*Corresponding author: frank.monahan@ucd.ie

**Short title**

**Effect of breed type and diet on bull beef quality**

## Abstract

In grass based beef production systems (PS), early maturing breed types (EM) may be preferable to late maturing breed types (LM) in achieving adequate fat cover. Biochemical and organoleptic characteristics of muscle from suckler bulls were investigated in EM and LM (n = 28/breed) assigned to one of two PS [*ad libitum* concentrates and grass silage to slaughter (C) or *ad libitum* silage plus 2 kg concentrate daily during winter followed by 99 days at pasture and then an indoor finishing period on C (GSPC)] in a 2 breed type × 2 PS factorial arrangement of treatments. Bulls were managed to have a common target carcass weight of 380 kg. Intramuscular fat (IMF) content was higher ( $P < 0.05$ ) for EM than LM, and for C than GSPC bulls. Collagen solubility was higher ( $P < 0.05$ ) for C than GSPC bulls. Lactate dehydrogenase (LDH) and phosphofructokinase activities were higher ( $P < 0.05$ ) for LM than EM. Isocitrate dehydrogenase activity and the Type I myosin heavy chain (MyHC) proportion were higher ( $P < 0.05$ ) for EM than LM. The LDH activity and the Type IIX MyHC proportion were higher ( $P < 0.05$ ) for C than GSPC bulls. Sensory ratings for tenderness and juiciness were higher ( $P < 0.01$ ) for beef from EM than LM while sensory ratings for tenderness, flavour liking and overall liking were higher ( $P < 0.001$ ) for C than for GSPC bulls. Differences in sensory quality were largely eliminated when adjusted for IMF. Overall, carcass fat scores, IMF and sensory scores were higher in EM than LM and in C than GSPC bulls but most differences in sensory quality could be attributed to differences in IMF.

**Key words:** beef, breed type, diet, sensory, intramuscular fat

## Implications

In countries like Ireland, where grazed grass is abundantly available, inclusion of grass silage followed by a period of grazed grass, prior to finishing on a high energy concentrate diet, decreases production costs in late maturing suckler bull production systems but the bulls may not meet the market-specific requirements in terms of carcass fat cover. It may be more appropriate, therefore, to rear early maturing breed types in such production systems as the bulls have higher carcass fat scores and marbling fat, and yield a more tender and juicier beef.

## Introduction

In Ireland, late maturing breed types (LM) account for 85% of the suckler beef herd while the remaining 15% are early maturing breed types (EM) (McGee, 2012). Traditionally, the male beef cattle population was dominated by steers, but more recently the proportion of bulls has increased as steers are less efficient in nutrient utilization than bulls when reared similarly (O'Riordan *et al.*, 2011). However, producing beef from suckler bulls, which usually involves provision of a high concentrate ration for a prolonged period, is usually less profitable because of the higher cost of concentrates compared to grass silage or grazed grass diets (Finneran *et al.*, 2011). Incorporating a grazing period prior to finishing on a concentrate diet has been shown to reduce the production costs of LM suckler bulls (O'Riordan *et al.*, 2011) with little impact on eating quality of the beef (Mezgebo *et al.*, 2016).

However, while it is economically viable to incorporate a grazing period in the LM suckler bull production system (PS), the bulls may not meet the market requirements in terms of adequate carcass fat cover at a particular carcass weight (O'Riordan *et al.*, 2011). Carcass fat cover and colour are important parameters for the beef industry as they influence the quality and consumer acceptability of beef (Moloney and Richardson, 2013). Even though LM predominate in the suckler herds in Ireland, EM may be more suitable for a grass-based PS because when managed to a particular slaughter weight and/or age, EM have a higher genetic potential to deposit fat than LM (Keane, 2011).

Recently, the influence, on beef quality characteristics, of incorporating a grazing period prior to indoor finishing on a concentrate diet in the LM suckler bull PS was evaluated

68 (Mezgebo *et al.*, 2016). However, to our knowledge, little is known about the effect of  
69 incorporating a grazing period in EM suckler bull PS on the quality of the beef.  
70 Therefore, the aim of this study was to determine the influences of breed maturity and  
71 inclusion of a period of grazed grass in a suckler bull PS on the compositional,  
72 biochemical and organoleptic characteristics of beef. It was hypothesised that LM could  
73 be replaced by EM, to achieve adequate fat cover and product quality specifications, in  
74 a suckler bull beef PS.

## Materials and methods

### *Animals and management*

As part of a larger study described by Marren *et al.* (2013), 28 spring-born (mean birth date 30 March) EM (Aberdeen Angus and Hereford sired calves) and 28 spring-born (mean birth date 8 March) LM (Charolais and Limousin sired calves) weaned suckler bulls were purchased at livestock markets in Ireland at approximately 8 months of age, acclimatised to slatted floor accommodation and offered grass silage *ad libitum* plus 2 kg/head/day of a barley-based concentrate. Bulls were randomly assigned (1 December) within breed maturity to a two breed types (B) × two PS factorial arrangement of treatments, balanced for sire breed and initial weight. The two PS were: (1) *ad libitum* concentrates (870 g/kg rolled barley, 60 g/kg soya bean meal, 50 g/kg molasses and 20 g/kg minerals/vitamins) plus *ad libitum* grass silage (GS) (dry matter digestibility 700 g/kg) (C), and (2) GS plus 2 kg concentrate daily during the winter (123 day duration) followed by 99 days at pasture and then an indoor finishing period on C (GSPC). Bulls were slaughtered at a commercial slaughter plant (Kepak Group, Clonee, Co. Meath, Ireland) on reaching a mean live weight estimated to achieve a target carcass weight of 380 kg. The study was carried out under license from the Irish Government and with the approval of Teagasc, the Agricultural and Food Development Authority.

### *Carcass grading and muscle tissue collection*

Post slaughter, carcasses were weighed and graded for conformation according to the EU Beef Carcass Classification Scheme as described in Mezgebo *et al.* (2016). At 1 h post-slaughter, a sample (ca. 20 g) of *longissimus thoracis* (LT) muscle tissue was taken (from 9<sup>th</sup> rib), snap frozen in liquid nitrogen and maintained at -80°C for muscle metabolic enzyme activity and muscle typing analyses.

### *Muscle pH and temperature measurement*

Muscle pH was measured at 2, 3.5, 5 and 48 h post-mortem by making a scalpel incision in the muscle at the 10<sup>th</sup> rib and inserting a glass electrode (Model EC-2010-06, Amagruss Electrodes Ltd., Westport, Co. Mayo, Ireland) attached to a portable pH meter (Model no. 250A, Orion Research Inc., Boston, MA) approximately 4.0 cm into

the muscle. The temperature was recorded simultaneously and used to make a temperature compensated pH measurement.

#### *Fat and muscle colour measurements*

A detailed procedure is given in Mezgebo *et al.* (2016). Briefly, at 48 h post-mortem, carcasses were cut at the 5/6<sup>th</sup> rib interface prior to subcutaneous fat and muscle colour measurements. Subcutaneous fat colour (i.e. *L*, *a*, *b* colour coordinates) was measured using a Miniscan XE Plus (Hunter Associates Laboratory Inc., Reston, VA, USA) at two positions: (1) the lower round/rump region and (2) 13<sup>th</sup> rib region. Chroma/saturation (*C*) and hue angle (*h*<sup>°</sup>) values were calculated from the '*a*' and '*b*' values. For muscle colour measurement, the cut surface of the muscle was first allowed to bloom for 1 h. Muscle colour grade was also subjectively assessed on the chilled carcass using Meat Standards Australia colour sticks (Anon, 2005). A portion of LT muscle (13 cm in length, from the 10<sup>th</sup> rib region) was excised, vacuum packed, aged for 14 days at 2°C, and finally frozen and stored at -18°C prior to compositional, collagen and sensory analysis.

#### *Proximate composition, collagen content and sensory analyses*

Moisture, intramuscular fat (IMF) and protein contents of the LT muscle were determined using the SMART System 5 microwave moisture drying oven, NMR SMART Trac rapid fat analyser (CEM Corporation, Matthews, NC, USA) and LECO FP328 (LECO Corp., St. Joseph, MI, USA) protein analyser, respectively (AOAC, 1990). Collagen content (i.e. total and soluble) was determined by quantitative determination of hydroxyproline by a colorimetric reaction (Kolar, 1990). Sensory analysis was carried out using a 10-person trained taste panel who had been selected for their sensory acuity, a detailed procedure is given in Mezgebo *et al.* (2016).

#### *Muscle metabolic enzyme activity and muscle contractile and metabolic type*

Glycolytic enzyme activities (lactate dehydrogenase (LDH) and phosphofructokinase (PFK)) and oxidative enzyme activities (isocitrate dehydrogenase (ICDH), citrate synthase (CS) and cytochrome c oxidase (COX)) were quantified spectrophotometrically according to Jurie *et al.* (2006). Muscle typing was assessed by determination of relative proportions of myosin heavy chains (MyHC) isoforms types I,

134 IIA and IIX using high-resolution mini-gel electrophoresis as described by Picard *et al.*  
135 (2011).

136 *Statistical analysis*

137 Data were subjected to analysis of variance using the General Linear Model procedure  
138 of SPSS (IBM SPSS Statistics Version 20) where the B, PS and their interaction were  
139 regarded as fixed factors. For data relating to sensory analysis, assessor and session  
140 effects were also included as fixed factors. The sensory data were also analysed using  
141 IMF as an overall linear covariate. Means were considered significant at  $P < 0.05$ .



## Results

### *Production and carcass traits*

Production, carcass and subcutaneous fat colour data are presented in Table 1. There was an interaction ( $P < 0.001$ ) between B and PS with respect to age at slaughter. Thus for C bulls, age at slaughter was higher for EM than for LM, but for GSPC bulls, age at slaughter was similar for EM and LM. The ADG indoor (i.e. during finishing on the concentrate diet) was lower ( $P < 0.001$ ) for C than for GSPC. There was an interaction ( $P < 0.05$ ) between B and PS with respect to ADG overall. Thus for C bulls, ADG overall was lower for EM than for LM, but for GSPC bulls, ADG overall was similar for EM and LM. Conformation score was lower ( $P < 0.001$ ) for EM than for LM. There was an interaction ( $P < 0.001$ ) between B and PS with respect to fat score. Thus for C bulls, fat score was similar for EM and LM, but for GSPC bulls, fat score was higher for EM than for LM. Subcutaneous fat 'L' and 'b' values were higher ( $P < 0.05$ ) for EM than for LM, and for C than for GSPC. 'h<sup>o</sup>' value was higher ( $P < 0.05$ ) for C than for GSPC bulls.

### *Muscle pH, temperature, colour, proximate composition and collagen data*

Muscle pH, temperature, colour, proximate composition and collagen data are presented in Table 2. At 2 h post-mortem, muscle pH was higher for EM than for LM ( $P < 0.001$ ), and for C than for GSPC ( $P < 0.01$ ). There was an interaction ( $P < 0.05$ ) between B and PS with respect to pH at 3.5 h post-mortem. Thus for EM, pH at 3.5 h was higher for C than for GSPC, but for LM, pH at 3.5 h was similar for C and GSPC. There was an interaction ( $P < 0.01$ ) between B and PS with respect to pH at 5 h post-mortem. Thus for C bulls, pH at 5 h was higher for EM than for LM, but for GSPC bulls, pH at 5 h was lower for EM than for LM. There was an interaction ( $P < 0.05$ ) between B and PS with respect to ultimate pH (pH<sub>u</sub>), i.e. 48 h post-mortem. Thus for C bulls, pH<sub>u</sub> was similar for EM and LM, but for GSPC bulls, pH<sub>u</sub> was higher for EM than for LM. There was an interaction ( $P < 0.001$ ) between B and PS with respect to muscle temperature at 2 h post-mortem. Thus for C bulls, muscle temperature at 2 h was lower for EM than for LM, but for GSPC bulls, muscle temperature at 2 h was higher for EM than for LM. At 3.5 h post-mortem, muscle temperature was higher ( $P < 0.001$ ) for EM than for LM. There was an interaction between B and PS with respect to muscle

temperature at 5 h post-mortem. Thus for C bulls, muscle temperature at 5 h post-mortem was similar for EM and LM, but for GSPC bulls, muscle temperature at 5 h post-mortem was higher ( $P < 0.01$ ) for EM than for LM. At 48 h post-mortem, muscle temperature was higher ( $P < 0.001$ ) for C than for GSPC.

For muscle colour, 'L' value was higher ( $P < 0.001$ ) for C than for GSPC, and 'a' value was higher ( $P < 0.001$ ) for GSPC than for C. There was an interaction ( $P < 0.05$ ) between B and PS with respect to 'b', 'C' and 'h<sup>o</sup>' values. Thus for C bulls, 'b', 'C' and 'h<sup>o</sup>' values were lower for EM than for LM, but for GSPC bulls, 'b', 'C' and 'h<sup>o</sup>' values were similar for EM and LM. Muscle colour grade was higher ( $P < 0.05$ ) for GSPC than for C. The IMF content was higher ( $P < 0.001$ ) for EM than for LM, and for C than for GSPC. Moisture content was higher for LM than for EM ( $P < 0.001$ ), and for GSPC than for C ( $P < 0.05$ ). Total collagen was higher ( $P < 0.05$ ) for EM than for LM. There was an interaction ( $P < 0.05$ ) between B and PS with respect to percentage of soluble collagen. Thus for C bulls, percentage of soluble collagen was higher for EM than for LM, but for GSPC bulls, percentage of soluble collagen was similar for EM and LM.

#### *Muscle metabolic enzyme activity and muscle contractile and metabolic type*

Muscle metabolic enzyme activity and MyHC proportion data are presented in Table 3. When enzyme activity was expressed as  $\mu\text{mol}/\text{min}$  per g of tissue, LDH activity was higher for LM than for EM ( $P < 0.001$ ), and for C than for GSPC bulls ( $P < 0.05$ ); PFK activity was higher ( $P < 0.05$ ) for LM than for EM; ICDH activity was higher ( $P < 0.01$ ) for EM than for LM and COX activity tended to be higher ( $P < 0.07$ ) for EM than for LM. When enzyme activity was expressed as  $\mu\text{mol}/\text{min}$  per g of protein, similar trends were observed although significance ( $P < 0.05$ ) was only reached in the case of the breed type effects on LDH and ICDH activities. Type I MyHC proportion was higher ( $P < 0.001$ ) for EM than for LM. Type IIX MyHC proportion was higher ( $P < 0.05$ ) for C than for GSPC.

#### *Sensory characteristics*

Muscle sensory data are presented in Table 4. Tenderness, flavour liking and overall liking were higher ( $P < 0.001$ ) for C than for GSPC. Tenderness and juiciness were higher ( $P < 0.01$ ) for EM than for LM. Ease of cutting ( $P < 0.001$ ) and cleanness of cut

203 ( $P < 0.05$ ) were higher for C than for GSPC. Clean cut was higher ( $P < 0.05$ ) for EM  
204 than for LM. Toughness (both during in-bite and eating) was higher for GSPC than for C  
205 ( $P < 0.001$ ), and for LM than EM ( $P < 0.05$ ). Juiciness (during in-bite) was higher for C  
206 than for GSPC ( $P < 0.01$ ), and for EM than for LM ( $P < 0.001$ ). Sponginess was higher  
207 ( $P < 0.001$ ) for C than for GSPC. Moisture, greasiness and pulpiness (both during  
208 eating and residual), dissolubility, ease of swallow and mouthfeel were higher ( $P < 0.05$ )  
209 for C than GSPC, and for EM than LM. Chewiness, fibrousness and residual particles  
210 were higher ( $P < 0.05$ ) for GSPC than for C. When the sensory data were analysed  
211 using IMF as a covariate, only beefy flavour was lower ( $P < 0.05$ ) and moisture and  
212 pulpiness (during eating) were higher ( $P < 0.05$ ) for EM than for LM (mean values of  
213 4.39 vs 4.59, 50.8 vs 46.9 and 55.8 vs 52.0 for beefy flavour, moisture and pulpiness  
214 respectively). Ease of swallow was higher ( $P < 0.05$ ) for C than GSPC (mean values of  
215 60.0 vs 54.7).

## Discussion

The bulls were slaughtered on reaching a mean group live weight estimated to achieve a target carcass weight of 380 kg which is required by some markets (Bord Bia, 2011). To reach the same target carcass weight, the LM bulls reared in the C PS grew faster generally (i.e. higher ADG overall), reached the desired live weight earlier and therefore were slaughtered at a younger age compared to that of EM bulls on the same PS. This confirms that LM are better converters of a high energy diet to carcass weight (Keane, 2011). However, when reared on the GSPC system, both breed types grew at a slower rate overall and took longer to reach the target live weight. Prior to slaughter (i.e. finishing period), the GSPC bulls grew faster compared to C bulls. The higher growth rate prior to slaughter for the GSPC bulls suggests compensatory growth during the indoor period as they had received a low energy diets (i.e. grass at pasture) prior to the finishing period compared to C bulls (Hornick *et al.*, 2000).

When managed to the same carcass weight, carcasses from LM are characterised by having relatively more muscle and less fat compared to carcasses from EM (O'Riordan *et al.*, 2011, Keane, 2011). In the present study, the better carcass conformation of the LM bulls compared to the EM bulls can be attributed to a higher degree of muscularity in the LM carcasses. Fat score, which is a measure of subcutaneous fat thickness or degree of finish, was similar between EM and LM in the C group possibly because of rapid growth due to the high energy diet of the C diet. However, in the GSPC bulls, carcasses of the LM were leaner even though both breed types were finished on the same concentrate diet. In this case, it appears that during the concentrate finishing period the LM were physiologically 'younger' and therefore were depositing less fat than the physiologically 'older' EM (Warriss, 2010). With regard to subcutaneous fat colour, the higher lightness of fat from EM compared to LM, and for C compared to GSPC bulls may be attributed to the higher fat scores (i.e. subcutaneous fat thickness over the muscle) of the carcasses of EM and C groups. Fat yellowness, often associated with grass diets due to accumulation of carotenoids (Dunne *et al.*, 2006), and reported to negatively influence consumer acceptability (Cornforth, 1994), was unexpectedly higher for C bulls compared to GSPC bulls. However, although differences in fat yellowness due to B and PS were significant ( $P < 0.05$ ), values were numerically quite similar,

suggesting that these colour differences would probably not be perceived by consumers. In the case of PS this may be attributed to the similarity in diets in the immediate pre-slaughter period.

The extent of post-mortem pH decline in a muscle depends on the glycogen concentration at slaughter which in turn depends on the animal's physical activity, nutrition and/or stress prior to slaughter (Klont and Lambooy, 1995; Warriss, 2010). In the present study, the influence of pre-slaughter physical activity and stress on muscle glycogen level would likely be minimal as the bulls were finished indoors and therefore were familiar with pre-slaughter handling; in addition the animals were carefully managed during transport and lairage. However, early post-mortem (i.e. 2, 3.5 and 5 h), a lower pH was recorded in the muscle from GSPC bulls compared to C bulls. This may be related to the higher growth rate of GSPC bulls during the finishing period compared to C bulls, whereby muscle is believed to become more glycolytic during periods of compensatory growth (Brandstetter et al., 1998). Similarly, a higher pH<sub>u</sub> (i.e. pH at 48 h post-mortem) was recorded in the muscle from EM breed types than LM breed types; however, there was an interaction between B and PS whereby the difference was observed in GSPC bulls and not in C bulls. The lower pH<sub>u</sub> for LM GSPC bulls could possibly reflect a higher muscle glycolytic potential as LM breed types are often characterised by an accelerated lean tissue growth compared to EM breed types when reared similarly (Hocquette *et al.*, 1998), in this case to a similar carcass weight. In agreement, glycolytic enzyme activity (LDH and PFK) were higher in muscle from LM breed types, as discussed further below. The higher muscle temperature at 3.5 h post-mortem for EM than LM bulls, and at 5 and 48 h post-mortem for C than GSPC bulls is most probably related to the carcass fat score as carcasses from EM and C groups had higher fatness scores than LM and GSPC groups, respectively. This is due to the fact that carcasses with a thicker fat cover cool more slowly than carcasses with a thinner fat cover (Warriss, 2010).

With regard to muscle colour, the lower lightness, higher redness, colour saturation and muscle colour grade (i.e. the higher the value, the darker the muscle) for the GSPC bulls could be explained by the higher age at slaughter (15.9 vs 18.5 months for C vs GSPC, respectively) as muscle tissue becomes darker and redder with increasing

278 slaughter age (Dunne *et al.*, 2006). The lower proportion of Type IIX MyHC, a  
279 characteristic of white muscles, for the GSPC bulls could also be responsible for the  
280 lower lightness of their LT muscle (Henckel *et al.*, 1997). The darker muscle from GSPC  
281 compared to C bulls could also be related to the physical activity during the pasture  
282 feeding period (Priolo *et al.*, 2001). However, it should be mentioned that the post-  
283 mortem pH profile of each muscle was within an acceptable pH range (Warriss, 2010),  
284 and thus meat from either group could not be considered to have experienced the 'dark  
285 cutting beef' condition.

286 The higher IMF content for EM compared to LM may be related to the intrinsic variations  
287 in the physiology of the animals (Oddy *et al.*, 2001) whereby at a similar live weight, the  
288 EM bulls were physiologically 'older' and therefore were depositing more IMF than the  
289 LM bulls, which were 'younger' physiologically, and therefore were depositing less IMF.  
290 The higher IMF content for C bulls reflects the higher energy content of the concentrate  
291 diet through out their life (Oddy *et al.*, 2001). The lower collagen solubility for GSPC  
292 bulls may be related to the greater age at slaughter (Blanco *et al.*, 2013) and lower IMF  
293 content (Nishimura, 2015) as an increase in slaughter age increases the proportion of  
294 mature collagen crosslinks which in turn leads to a decrease in solubility of the collagen  
295 (McCormick, 1994).

296 The higher glycolytic enzyme activities (LDH and PFK) for LM could be related to the  
297 higher overall growth rate of these bulls as an increase in growth rate early in life (i.e.  
298 period of rapid growth from one to 12 months) and further growing stage until sexual  
299 maturity is associated with an increase in muscle LDH activity (i.e. glycolytic  
300 metabolism) and a decrease in ICDH activity (i.e. oxidative metabolism) (Jurie *et al.*,  
301 1995). A similar explanation could be offered for the tendency towards lower oxidative  
302 enzyme activities ( $P < 0.07$ ) of ICDH and COX, marker enzymes for tricarboxylic acid  
303 cycle and mitochondrial electron transport respectively, and lower proportion of slow  
304 twitch Type I oxidative MyHC in the muscle from LM. The higher LDH activity (per g of  
305 tissue) and proportion of Type IIX (fast twitch glycolytic) MyHC for C bulls could be  
306 explained by the higher overall growth rate which is mainly attributed to the continued  
307 provision of concentrate diet which in turn results in a more glycolytic muscular  
308 metabolism (Brandstetter *et al.*, 1998, Cassar-Malek *et al.*, 2004). In addition, such

309 higher glycolytic metabolism in muscle could also be associated with the longer  
310 concentrate finishing period of the C group compared to GSPC group (i.e. concentrate  
311 finishing period of 98 and 71 d for GSPC and 258 and 201 d for C bulls of EM and LM  
312 respectively). Even though grazing on pasture is associated with an increase in  
313 oxidative metabolism of muscle mainly due to higher physical activity (Therkildsen *et al.*,  
314 1998), the C and GSPC groups had similar oxidative enzyme activities. However, this  
315 was not unexpected as all bulls were finished indoors on the same concentrate diets for  
316 at least 71 days. In the present study, the fast twitch Type IIB glycolytic muscle MyHC  
317 was expressed in only 6 bulls (1 in EM of C, none in EM of GSPC, 1 in LM of C and 4 in  
318 LM of GSPC bulls, data not shown) in contrast to a study by Picard and Cassar-Malek  
319 (2009) in a Blonde d'Aquitaine (a French beef breed) in which Type IIB MyHC was  
320 usually identified.

321 The effect of PS on sensory characteristics was in agreement with Mezgebo *et al.*  
322 (2016). The higher tenderness scores for C bulls may be related to their younger age at  
323 slaughter (Bures and Barton, 2012), higher IMF (Thompson, 2004) and collagen  
324 solubility (Cross *et al.*, 1973). A similar explanation could be given for the higher  
325 sensory ratings for ease of cutting, cleanness of cut, juiciness (in-bite), sponginess,  
326 moisture, greasiness, pulpiness, dissolubility, ease of swallow and mouthfeel, and lower  
327 ratings in toughness, chewiness, fibrousness and residual particles for C bulls  
328 compared to GSPC bulls. The contribution of IMF to these differences was shown by  
329 the lack of significant differences in sensory ratings (except for ease of swallow)  
330 between PS when the data were adjusted for IMF. In addition, the higher LDH activity  
331 for the C bulls compared to GSPC bulls could also be linked to the higher tenderness  
332 ratings of the C bulls, as an increase in glycolytic characteristics of a muscle often leads  
333 to an increase in eating quality of meat mainly by accelerating the post-mortem  
334 tenderization process of the muscle (Maltin *et al.*, 2001). The sensory analysis also  
335 showed that the sensory data ratings were internally consistent, especially for  
336 tenderness, i.e. higher tenderness score (during the basic taste) was consistent with the  
337 lower toughness scores (both during in-bite and eating). Even though all bulls were  
338 finished indoors, the lower flavour liking and overall liking ratings of beef from GSPC  
339 bulls could possibly be associated with the inclusion of grass diet prior to the finishing

340 period as beef from pasture based systems is often reported to be less preferred by  
341 consumers (Griebenow *et al.*, 1997).

342 The higher sensory ratings in tenderness and juiciness, and associated higher scores in  
343 cleanness of cut, moisture, greasiness, pulpiness, dissolubility, ease of swallow and  
344 mouthfeel, and lower scores in toughness for EM could be related to their higher  
345 carcass fat cover and IMF content compared to LM. Similar findings were reported by  
346 Sinclair *et al.* (2001) in beef from Aberdeen-Angus and Charolais breeds. In the current  
347 study, beef from LM was rated to be lower in tenderness, juiciness and related sensory  
348 quality attributes compared to beef from EM even though the LM were younger at  
349 slaughter. In addition, LM muscle had higher glycolytic (LDH and PFK) and lower  
350 oxidative (ICDH and COX) metabolic enzyme activities and lower Type I MyHC  
351 proportion than EM, and an increase in glycolytic (Maltin *et al.*, 2001) and decrease in  
352 oxidative (Monin and Ouali, 1991) characteristics of a muscle can lead to superior  
353 eating quality in meat. When IMF was included as a covariate in the sensory data  
354 analysis, most of the observed differences disappeared, confirming that IMF content  
355 was the major contributor to differences in meat tenderness and juiciness between EM  
356 and LM breeds (Sinclair *et al.*, 2001).



## **Conclusion**

When managed to a similar carcass weight EM were older at slaughter, had higher carcass fat scores and IMF content and produced beef that was rated more tender and juicier by trained sensory panellists than LM. Furthermore, C bulls were younger at slaughter, had higher carcass fat scores, IMF and soluble collagen content and produced beef rated more highly by a trained sensory panel than GSPC bulls. While variations in sensory characteristics due to breed maturity and dietary inclusion of grass silage followed by pasture exist, IMF contributed to much of the variation and it remains to be established whether or not the differences would be perceptible to untrained consumers.

## **Acknowledgements**

The senior author acknowledges receipt of a Teagasc Walsh Fellowship. The contributions of Dr Alan Kelly (UCD), Mr Declan Marren (Teagasc Animal and Grassland Research and Innovation Centre, Grange) and Ms Karen Hussey (Teagasc Ashtown Food Research Centre) are also acknowledged. The assistance of staff and management at Kepak Group, Clonee, Co. Meath who facilitated sample collection and carcass measurement is greatly appreciated.

## References

- Anon 2005. Handbook of Australian meat, 7th edition. AUS-MEAT Limited, Brisbane, Queensland, Australia.
- Association of Official Analytical Chemists (AOAC) 1990. Moisture and fat in meat and poultry products. Official Methods 985.14 and 985.26 (ed. P Cunniff), pp. 931-932. AOAC International, Arlington, VA, United States.
- Blanco M, Jurie C, Micol D, Agabriel J, Picard B and Garcia-Launay F 2013. Impact of animal and management factors on collagen characteristics in beef: a meta-analysis approach. *Animal* 7, 1208-1218.
- Bord Bia 2011. Bord Bia - Irish Food Board Report, Dublin, Ireland.
- Brandstetter AM, Picard B and Geay G 1998. Muscle fibre characteristics in four muscles of growing male cattle: Effect of castration and feeding level. *Livestock Production Science* 53, 25–36.
- Bures D and Barton L 2012. Growth performance, carcass traits and meat quality of bulls and heifers slaughtered at different ages. *Czech Journal of Animal Science* 57, 34-43.
- Cassar-Malek I, Hocquette JF, Jurie C, Listrat A, Jailler R, Bauchart D, Briand Y and Picard B 2004. Muscle-specific metabolic, histochemical and biochemical responses to a nutritionally induced discontinuous growth path. *Journal of Animal Science* 79, 49–59.
- Cornforth D 1994. Colour – its basis and importance. In *Quality attributes and their measurement in meat, poultry and fish products – Advances in Meat Research Series* (ed. AM Pearson and TR Dutson), Volume 9, pp. 34–78. Chapman and Hall, London, UK.
- Cross HR, Carpenter ZL and Smith GC 1973. Effects of intramuscular collagen and elastin on bovine muscle tenderness. *Journal of Food Science* 38, 998-1003.
- Dunne PG, O'Mara FP, Monahan FJ and Moloney AP 2006. Changes in colour characteristics and pigmentation of subcutaneous adipose tissue and *M. longissimus dorsi* of heifers fed grass, grass silage or concentrate based diets. *Meat Science* 74, 231-241.
- Finneran E, Crosson P, O'Kiely P, Shalloo L, Forristal D and Wallace M 2011. Stochastic simulation of the cost of home-produced feeds for ruminant livestock systems. *Journal of Agricultural Science* 150, 123-139.

404 Griebenow RL, Martz FA and Morrow RE 1997. Forage based beef finishing systems: A review.  
 405 Journal of Production Agriculture 10, 84-91.

406 Henckel P, Oksbjerg N, Erlandsen E, Bartongade P and Bejerholm C 1997. Histo- and  
 407 biochemical characteristics of the longissimus dorsi muscle in pigs and their relationships to  
 408 performance and meat quality. Meat Science 47, 311–321.

409 Hocquette J, Ortigues-Marty I, Pethick D, Herpin P and Fernandez X 1998. Nutritional and  
 410 hormonal regulation of energy metabolism in skeletal muscles of meat-producing animals.  
 411 Livestock Production Science 56, 115-143.

412 Hornick JL, Van Eenaeme C, Gérard O, Dufrasne I and Istasse L 2000. Mechanisms of reduced  
 413 and compensatory growth. Domestic Animal Endocrinology 19, 121-132.

414 Immonen K, Schaefer DM, Puolanne E, Kauffman RG and Nordheim EV 2000. The relative  
 415 effect of dietary energy density on repleted and resting muscle glycogen concentrations. Meat  
 416 Science 54, 155-162.

417 Jurie C, Robelin J, Picard B and Geay Y 1995. Post-natal changes in the biological  
 418 characteristics of Semitendinosus muscle in male Limousin cattle. Meat Science 41, 125-135.

419 Jurie C, Ortigues-Marty I, Picard B, Micol D and Hocquette J 2006. The separate effects of the  
 420 nature of diet and grazing mobility on metabolic potential of muscles from Charolais steers.  
 421 Livestock Science 104, 182-192.

422 Keane MG 2011. Ranking of sire breeds and beef cross breeding of dairy and beef cows.  
 423 Teagasc Occasional Series No. 9, March 1. Grange Beef Research Centre, Co. Meath, Ireland.  
 424 [http://t-stor.teagasc.ie/bitstream/11019/810/1/Ranking\\_of\\_sire\\_breeds.pdf](http://t-stor.teagasc.ie/bitstream/11019/810/1/Ranking_of_sire_breeds.pdf)

425 Klont RE and Lambooy E 1995. Effects of preslaughter muscle exercise on muscle metabolism  
 426 and meat quality studied in anesthetized pigs of different halothane genotypes. Journal of  
 427 Animal Science 73, 108-117.

428 Kolar K 1990. Colorimetric determination of hydroxyproline as measure of collagen content in  
 429 meat and meat products: NMKL collaborative study. Journal-Association of Official Analytical  
 430 Chemists 73, 54-57.

431 Maltin CA, Lobley GE, Grant CM, Miller LA, Kyle DJ, Horgan GW, Matthews KR and Sinclair KD  
 432 2001. Factors influencing beef eating quality: Effects of nutritional regimen and genotype on  
 433 muscle fibre characteristics. *Animal Science* 72, 279–287.

434 Marren D, McGee M, Moloney AP, Kelly A and O’Riordan EG 2013. Comparison of early- and  
 435 late-maturing suckler-bred bulls on contrasting production systems, slaughtered at three  
 436 carcass weights. *Proceedings of Agricultural Research Forum*, pp. 72. Tullamore, Co. Offaly,  
 437 Ireland.

438 McCormick RJ 1994. The flexibility of the collagen compartment of muscle. *Meat Science* 36,  
 439 79-91.

440 McGee M 2012. Review of Irish suckler cow types: Research perspective. In “Setting a New  
 441 Direction for Suckler Cow Breeding”. *Proceedings of the Teagasc-ICBF Suckler Cow Breeding*  
 442 *Conference*, Tullamore, Ireland, 11 October 2012, p1–19.

443 Mezgebo GB, Moloney AP, O’Riordan EG, McGee M, Richardson IR and Monahan FJ 2016.  
 444 Comparison of organoleptic quality and composition of beef from suckler bulls from different  
 445 production systems. *Animal*. doi:10.1017/S1751731116001944.

446 Moloney AP and Richardson IR 2013. Sensory characteristics of muscle from Angus x  
 447 Holstein/Friesian and Belgian Blue x Holstein/Friesian heifers slaughtered at two ages.  
 448 *Proceedings of Agricultural Research Forum*, pp. 32. Tullamore, Co. Offaly, Ireland.

449 Monin G and Ouali A 1991. Muscle differentiation and meat quality. In *Developments in meat*  
 450 *science* (ed. RA Lawrie), pp. 89–157. Elsevier Applied Science, London, UK.

451 Nishimura T 2015. Role of extracellular matrix in development of skeletal muscle and  
 452 postmortem aging of meat. *Meat Science* 109, 48-55.

453 O’Riordan EG, Crosson P and McGee M 2011. Finishing male cattle from the beef suckler herd.  
 454 *Irish Grassland Association Journal* 45, 131-146.

455 Oddy VH, Harper GS, Greenwood PL and McDonagh MB 2001. Nutritional and developmental  
 456 effects on the intrinsic properties of muscles as they relate to the eating quality of beef.  
 457 *Australian Journal of Experimental Agriculture* 41, 921–942.

458 Picard B and Cassar-Malek I 2009. Evidence for expression of IIb myosin heavy chain isoform  
 459 in some skeletal muscles of Blonde d’Aquitaine bulls. *Meat Science* 82, 30-36.

460 Picard B, Barboiron C, Chadeyron D and Jurie C 2011. Protocol for high- resolution  
 461 electrophoresis separation of myosin heavy chain isoforms in bovine skeletal muscle.  
 462 Electrophoresis 32, 1804-1806.

463 Priolo A, Micol D and Agabriel J 2001. Effects of grass feeding systems on ruminant meat  
 464 colour and flavour. A review. Journal of Animal Research 50, 185-200.

465 Sinclair K, Lobley G, Horgan G, Kyle D, Porter A, Matthews K, Warkup C and Maltin C 2001.  
 466 Factors influencing beef eating quality. 1. Effects of nutritional regimen and genotype on  
 467 organoleptic properties and instrumental texture. Animal Science 72, 269-278.

468 Therkildsen M, Vestergaard M, Jensen LR, Andersen HR and Sejrsen K 1998. Effect of feeding  
 469 level, grazing and fnishing on growth and carcass quality of young Friesian bulls. Acta Agri-  
 470 culturae Scandinavica, Section A. Animal Science 48, 193-201.

471 Thompson J 2004. The effects of marbling on flavour and juiciness scores of cooked beef, after  
 472 adjusting to a constant tenderness. Australian Journal of Experimental Agriculture 44, 645-652.

473 Warriss PD 2010. Meat Science: An Introductory Text, 2nd edition. CABI Publishing, London,  
 474 UK.

**Table 1** Production, carcass and subcutaneous fat colour data of bulls from two breed types (B) (EM = early maturing, LM = late maturing), raised on two production systems (PS) (C = concentrate, GSPC = grass silage followed by pasture and then concentrate)

	B	EM		LM		s.e.m	Significance		
	PS	C	GSPC	C	GSPC		B	PS	B x PS
Finishing period (days) <sup>1</sup>		258	98	201	71				
Age at slaughter (months)		16.7 <sup>b</sup>	18.6 <sup>c</sup>	15.0 <sup>a</sup>	18.3 <sup>c</sup>	0.25	***	***	**
ADG <sup>2</sup> finishing (kg/day)		1.35	2.09	1.50	2.06	0.081		***	
ADG overall (kg/day)		1.38 <sup>b</sup>	1.09 <sup>a</sup>	1.58 <sup>c</sup>	1.10 <sup>a</sup>	0.042	*	***	*
Slaughter weight (kg)		681	704	667	693	14.1			
Carcass weight (kg)		375	385	379	387	9.1			
Conformation score <sup>3</sup>		8.3	8.7	9.9	9.7	0.36	***		
Fat score <sup>4</sup>		8.3 <sup>b</sup>	8.3 <sup>b</sup>	8.4 <sup>b</sup>	6.6 <sup>a</sup>	0.26	***	***	***
Fat colour <sup>5</sup>									
‘L’		72.4	68.9	68.6	64.5	0.66	***	***	
‘a’		9.1	9.5	8.8	9.3	0.50			
‘b’		16.9	15.6	15.6	15.4	0.32	*	*	
‘C’		19.3	18.3	17.9	18.0	0.46			
‘h°’		62.0	58.9	61.1	58.9	1.16		*	

<sup>1</sup> Days on *ad libitum* concentrates prior to slaughter

<sup>2</sup>Average daily live weight gain

<sup>3</sup>Conformation classes E<sup>+</sup> (highest) to P<sup>-</sup> (lowest), (E<sup>+</sup> is 15)

<sup>4</sup>Fat score classes 5<sup>+</sup> (highest) to 1<sup>-</sup> (lowest), (5<sup>+</sup> is 15)

<sup>5</sup>Subcutaneous fat colour: ‘L’ = lightness, 0 (black) to 100 (white); ‘a’ = redness, +a (red) to –a (green); ‘b’ = yellowness, +b (yellow) to –b (blue); ‘C’ = chroma, higher ‘C’ values higher colour saturation; ‘h°’ = hue, 0/360° is red, 90° is yellow, 180° is green and 270° is blue colour

<sup>a, b, c</sup> means within rows (where interaction exists), assigned different superscripts differ significantly (*P* < 0.05)

\**P* < 0.05, \*\**P* < 0.01, \*\*\**P* < 0.001

**Table 2** Post-mortem pH and temperature, colour, proximate composition and collagen content of longissimus thoracis muscle from bulls from two breed types (B) (EM = early maturing, LM = late maturing), raised on two production systems (PS) (C = concentrate, GSPC = grass silage followed by pasture and then concentrate)

B		EM		LM		Significance			
PS	C	GSPC	C	GSPC	s.e.m	B	PS	B x PS	
pH, post-mortem (h)									
2	6.59	6.47	6.45	6.28	0.054	***	**		
3.5	6.21 <sup>c</sup>	5.84 <sup>a</sup>	6.11 <sup>bc</sup>	5.97 <sup>ab</sup>	0.054		***	*	
5	6.03 <sup>c</sup>	5.67 <sup>a</sup>	5.87 <sup>b</sup>	5.85 <sup>b</sup>	0.056		***	**	
48	5.69 <sup>ab</sup>	5.74 <sup>b</sup>	5.68 <sup>ab</sup>	5.62 <sup>a</sup>	0.026	**		*	
Temperature, post-mortem (h)									
2	33.1 <sup>a</sup>	35.1 <sup>b</sup>	35.3 <sup>b</sup>	32.4 <sup>a</sup>	0.55			***	
3.5	29.1	29.5	28.1	27.3	0.48	***			
5	23.9 <sup>b</sup>	24.1 <sup>b</sup>	24.4 <sup>b</sup>	21.9 <sup>a</sup>	0.47		*	**	
48	3.90	3.25	3.66	3.09	0.154		***		
Muscle colour <sup>1</sup>									
‘L’	31.1	28.1	32.8	28.3	0.80		***		
‘a’	19.8	21.6	20.5	21.2	0.30		***		
‘b’	12.2 <sup>a</sup>	12.9 <sup>a</sup>	13.9 <sup>b</sup>	12.9 <sup>a</sup>	0.24	***		***	
‘C’	23.3 <sup>a</sup>	25.1 <sup>b</sup>	24.8 <sup>b</sup>	24.8 <sup>b</sup>	0.32		**	**	
‘h <sup>o</sup> ’	31.7 <sup>a</sup>	30.8 <sup>a</sup>	34.2 <sup>b</sup>	31.4 <sup>a</sup>	0.50	***	***	*	
Muscle colour grade <sup>2</sup>	3.07	3.29	2.57	3.21	0.172		*		
Proximate composition (g/kg)									
Intramuscular fat	55.2	27.7	26.2	10.2	3.94	***	***		
Moisture	720	738	747	749	4.8	***	*		
Protein	229	233	229	231	2.7				
Ash	10.5	12.0	11.2	11.3	0.59				
Collagen content									
Total collagen (mg/g)	4.06	4.21	3.86	3.87	0.126	*			
Soluble collagen (%)	13.4 <sup>b</sup>	8.3 <sup>a</sup>	9.4 <sup>a</sup>	9.4 <sup>a</sup>	0.79		***	***	

<sup>1</sup>Muscle colour: ‘L’ = lightness, 0 (black) to 100 (white); ‘a’ = redness, +a (red) to –a (green); ‘b’ = yellowness, +b (yellow) to –b (blue); ‘C’ = chroma, higher ‘C’ values higher colour saturation; ‘h<sup>o</sup>’ = hue, 0/360° is red, 90° is yellow, 180° is green and 270° is blue colour

<sup>2</sup>Muscle colour grades: 1 (extremely bright red) to 9 (extremely dark red)

<sup>a, b, c</sup>means within rows (where interaction exists), assigned different superscripts differ significantly ( $P < 0.05$ )

\* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$

499 **Table 3** Metabolic enzyme activity and myosin heavy chains (MyHC) proportion of  
 500 *longissimus thoracis muscle* from bulls from two breed types (B) (EM = early maturing,  
 501 LM = late maturing), raised on two production systems (PS) (C = concentrate, GSPC =  
 502 grass silage followed by pasture and then concentrate)

B		EM		LM		s.e.m	Significance		
PS	C	GSPC	C	GSPC	B		PS	B x PS	
Metabolic enzyme activity <sup>1</sup>									
μmol/min per g of tissue									
LDH	936	838	999	969	26.5	***	*		
PFK	101	96	112	112	6.8	*			
ICDH	1.17	1.33	1.01	1.02	0.085	**			
COX	17.0	18.3	15.1	15.2	1.33	0.07			
CS	5.27	5.37	5.34	4.58	0.463				
μmol/min per g of protein									
LDH	4908	4350	5007	5478	275.9	*			
PFK	527	498	559	636	45.0	0.06			
ICDH	6.14	6.90	5.12	5.68	0.483	*			
COX	89.3	94.7	75.7	87.1	8.14				
CS	27.7	27.9	27.1	26.2	2.86				
Protein (mg/g of tissue)	191	193	200	186	4.7				
MyHC <sup>2</sup> proportion (%)									
I	22.5	23.2	18.5	17.1	1.64	***			
IIA	45.1	48.8	38.6	46.8	3.35				
IIX	35.3	32.7	44.1	29.8	3.44		*		

503 <sup>1</sup>LDH: lactate dehydrogenase; PFK: phosphofructokinase; ICDH: isocitrate dehydrogenase;  
 504 COX: cytochrome c oxidase; CS: citrate synthase  
 505 <sup>2</sup>I: oxidative, IIA: oxido-glycolytic, IIX: glycolytic  
 506 \**P* < 0.05, \*\**P* < 0.01, \*\*\**P* < 0.001



**Table 4** Sensory characteristics of of longissimus thoracis muscle from bulls from two breed types (B) (EM = early maturing, LM = late maturing), raised on two production systems (PS) (C = concentrate, GSPC = grass silage followed by pasture and then concentrate)

B		EM		LM		s.e.m.	Significance		
PS	C	GSPC	C	GSPC	B		PS	B x PS	
Basic tastes, scale 1 (least) - 8 (most)									
Tenderness	4.81	4.50	4.63	4.20	0.093	**	***		
Juiciness	5.10	4.90	4.83	4.81	0.068	**			
Beefy flavour	4.54	4.41	4.55	4.51	0.060				
Abnormal flavour	2.30	2.50	2.30	2.42	0.074				
Flavour liking	5.45	5.02	5.46	5.10	0.081		***		
Overall liking	5.15	4.71	5.03	4.59	0.081		***		
Specific sensory indicators, scale 0 (nil) - 100 (extreme)									
On-cut									
Ease of cutting	55.7	49.6	53.5	46.7	1.34		***		
Cleanness of cut	59.2	56.8	56.6	53.9	1.20	*	*		
In-bite									
Toughness	43.1	48.8	45.5	54.9	1.35	**	***		
Crispness	25.3	26.1	24.3	25.6	1.08				
Juiciness	51.1	47.5	46.7	44.2	1.03	***	**		
Sponginess	29.9	26.9	28.6	25.5	0.87		***		
Eating									
Toughness	43.1	48.7	44.9	53.5	1.33	*	***		
Moisture	52.2	49.6	48.1	45.0	1.05	***	**		
Chewiness	40.9	47.1	42.6	49.0	1.42		***		
Greasiness	21.5	17.9	19.1	15.7	0.88	**	***		
Fibres	42.1	43.2	42.6	46.1	1.05		*		
Gristle	5.5	6.2	6.4	6.2	0.68				
Pulpy	57.5	54.9	52.7	50.2	1.08	***	**		
Dissolubility	51.5	46.3	49.6	43.0	1.31	*	***		
Residual									
Greasiness	21.5	18.2	18.4	15.4	0.93	**	***		
Ease of swallow	62.1	55.3	59.5	52.5	1.21	*	***		
Pulpy	56.7	54.4	51.9	48.2	1.12	***	**		
Particles	49.6	50.3	48.9	52.5	0.99		*		
Mouthfeel	57.0	54.5	52.2	49.8	0.99	***	*		

\**P* < 0.05, \*\**P* < 0.01, \*\*\**P* < 0.001